arXiv:1110.6167v1 [math.DS] 27 Oct 2011

# HOMOGENEOUS APPROXIMATION FOR FLOWS ON TRANSLATION SURFACES

JON CHAIKA

Given a translation surface $M$ with a distinguished horizontal direction, we denote the flow in direction $2\pi\theta$ to the horizontal by $F_\theta$. Let $d$ denote distance on $M$.

We say $M$ is a *Khinchin surface* if for every nonincreasing function $f : \mathbb{R}^+ \to \mathbb{R}^+$ such that $\int_C^\infty f(t)dt = \infty$ for any $C > 0$ we have that for almost every $\theta \in [0, 1)$ and $x \in M$, $d(F_\theta^t(x), x) < f(t)$ for arbitrarily large $t$.

**Theorem 1.** *All translation surfaces are Khinchin surfaces.*

*Corollary* 1. Let $\{a_i\}_{i=1}^\infty$ be a non-increasing sequence of positive real numbers with $\sum_{i=1}^{\infty} a_i = \infty$. For almost every IET $T$ and almost every point $x$ we have $\liminf_{n\to\infty} a_n^{-1}|T^n x - x| = 0$. That is, $|T^n x - x| < a_n$ for infinitely many $n$.

We were motivated by:

**Theorem 2.** *(Marchese [5]) Let $\{a_i\}_{i=1}^\infty$ be a decreasing sequence with divergent sum and with the additional property that $\{ia_i\}_{i=1}^\infty$ is decreasing. For almost every IET $T$*

$$\delta \in \bigcap_{n=1}^{\infty} \bigcup_{i=n}^{\infty} B(T^i(\delta'), a_i))$$

*where $\delta$ and $\delta'$ are any discontinuities of $T$.*

*Remark* 1. Neither Corollary 1 or Theorem 2 is stronger than the other. Corollary 1 allows more general sequences and treating almost every point. Marchese has recently adapted Theorem 2 to treat almost every point. Additionally, Corollary 1 lets us treat a one parameter family obtained by fixing any translation surface and considering varying the direction of flow and examining first return to a transversal on a fixed translation surface. By treating each translation surface, it applies to billiards in rational polygons [3]. It does not recover Theorem 2's result that all discontinuities approximate each other.

Let $M$ be a translation surface of genus $g$ and $s$ be the length of the shortest saddle connection on $M$. Let $\sigma = \frac{1}{2g-2}$. Let $S_\sigma(M)$ be the sequence of directions of periodic cylinders of volume at least $\sigma$ enumerated by increasing length. Let $\tilde{S}_\sigma(L) =$

$$\{(\theta, T) : \theta \text{ is the direction of a periodic cylinder of length } T < L \text{ and volume } \sigma\}.$$

**Proposition 1.** *(Vorobets)Let $m$ be the sum of the multiplicities of singularities. Let $c = 2^{2^{4m}}\sqrt{s}$. For large enough $L$ we have $\bigcup_{(\theta,T)\in\tilde{S}_\sigma(L)} B(\theta, \frac{c}{TL}) = S^1$.*

This is shown on [6, page 14].

The following proposition is a key result.

**Proposition 2.** *There exist* $C_1, C_2, C_3 > 0$ *such that for any interval* $J$ *and* $N > C_3$ *we have* $\lambda(\bigcup_{\theta \in \tilde{S}_\sigma(C_1N) \setminus \tilde{S}_\sigma(N)} B(\theta, \frac{1}{(C_1N)^2}) \cap J) > C_2\lambda(J) - \frac{2}{(C_1N)}^2$.

To prove this we establish a few other results.

**Lemma 1.** *If* $x$ *is in a periodic cylinder with direction* $\theta$*, length* $L$ *and area* $\sigma$ *then* $x$ *is not in a periodic cylinder of direction* $\phi$ *and length* $R$ *for any* $\phi \in B(\theta, \frac{1}{RL})$.

*Proof.* Notice that, $d(F_\theta^{t\cos(\theta-\theta')}(x), F_{\theta'}^t(x)) = |\sin(\theta-\theta')|t$ so long as no singularity lies on the geodesic connecting $F_\theta^s(x)$ to $F_{\theta'}^s(x)$ for $0 \leq s \leq t$. The cylinder must have circumference $\frac{\sigma}{L}$. If $x$ is in a periodic direction of length $R$ and direction $\phi$ then $F_\phi^s(x) = F_\phi^{s-R}(x)$ for all $s$. Choose $s$ so that $F_\phi^s(x)$ is just leaving the periodic cylinder in direction $\theta$ that $x$ is in. So $F_\phi^{s-t}(x)$ is in this periodic cylinder for all $t < \frac{\sigma}{L}|\csc(\theta-\phi)|$ and in particular $F_\phi^{s-t}(x) \neq F_\phi^s(x)$. So $R \geq \frac{\sigma}{L}|\csc(\theta-\phi)|$ which implies that $|\theta - \phi| \geq \frac{1}{RL}$. ☐

*Corollary* 2. For any interval $J$ we have $\lambda(\bigcup_{(\theta,T)\in\tilde{S}_\sigma(L)} B(\theta, \frac{1}{TL}) \cap J) \leq \sigma^{-1}\lambda(J)$.

*Proof.* If $(\theta, T) \in \tilde{S}_\sigma(L)$ then there exists a periodic cylinder $U$ in direction $\theta$ with length $T$. If $x \in U$ then $x$ is not in another periodic cylinder of length less than $L$ and direction $\phi \in B(\theta, \frac{1}{TL})$. So any direction in $J$ can be contained in at most $\sigma^{-1}$ of the $B(\theta', \frac{1}{LT})$ for $(\theta', L) \in \tilde{S}_\sigma(L)$. ☐

*Corollary* 3. There exists $C_5 > 0$ such that for all large enough $L$ we have

$$|\{(\theta, T) \in \tilde{S}_\sigma(C_5L) \setminus \tilde{S}_\sigma(L) : \theta \in J\}| > \frac{C_1}{2}\lambda(J)L^2.$$

*Proof.* Proposition 1 and Corollary 2 imply that

$$C_1\lambda(J) \leq \sum_{(\theta,T)\in\tilde{S}_\sigma(L), B(\theta,\frac{1}{TL})\cap J\neq\emptyset} \frac{2}{TL} \leq \sigma^{-1}(\lambda(J) + \frac{2}{L}).$$

Therefore for large enough $L$ we have $\sum_{(\theta,T)\in\tilde{S}_\sigma(L)} \frac{2}{T(3C_1^{-1}\sigma^{-1})L} < \frac{C_1\lambda(J)}{2}$. So

$$\sum_{(\theta,T)\in\tilde{S}_\sigma(3C_1^{-1}\sigma^{-1}L)\setminus\tilde{S}_\sigma(L), B(\theta,\frac{1}{TL})\cap J\neq\emptyset} \frac{2}{3C_1^{-1}\sigma^{-1}TL} \geq \frac{C_1\lambda(J)}{2}.$$

The corollary follows. ☐

*Proof of Proposition 2.* This is a consequence of Lemma 1 and the previous corollary because for fixed $k > 1$ and $(\theta, T) \in \tilde{S}_\sigma(kR) \setminus \tilde{S}_\sigma(R)$ we have $TkR$ is proportional to $(kR)^2$. ☐

We now state a useful theorem which follows from [1] and is phrased as in [2, Theorem 1].

**Theorem 3.** *If $\{x_i\}_{i=1}^{\infty} \subset [0,1]$ is a sequence with the property that there exists $c > 0$ such that for any interval $J$ we have*

$$\liminf_{N\to\infty} \lambda\Big(\bigcup_{i=1}^{N} B(x_i, s_i) \cap J\Big) > c\lambda(J)$$

*then for any non increasing sequence of positive real numbers $r_1 \geq r_2 \geq \ldots$ with $\sum a_i = \infty$ we have $\lambda(\bigcap_{n=1}^{\infty} \bigcup_{i=n}^{\infty} B(s_i, r_i)) = 1$.*

Observe that Proposition 2 shows that $S_\sigma(M)$ satisfies the hypothesis of this theorem. So by fact that the number of cylinders is quadratic in the length (that is, the $k^{th}$ element has length proportional to $\sqrt{k}$) we obtain the following corollary.

*Corollary* 4. If If $f$ is a non-increasing positive function such that $\int_C^{\infty} f(\sqrt{t})dt = \infty$ for any $C > 0$ then for almost every $\phi$ and any $L_0 > 0$ there exists $(\theta, T) \in S_\sigma(\infty)$ with $T > L_0$ and $|\phi - \theta| < f(T)$.

**Lemma 2.** *Let $\theta$ be the direction of a cylinder of length $L$ and volume $a$. If $|\theta - \phi| \leq \frac{\epsilon}{L}$ then for a set of points of measure at least $\frac{a}{4}$ we have $d(F_\phi^{L|\sec(\theta-\phi)|}x, x) < 2\epsilon$.*

*Proof.* $d(F_\theta^{t\cos(\theta-\theta')}(x), F_{\theta'}^{t}(x)) = |\sin(\theta - \theta')|t$ so long as no singularity lies on the geodesic connecting $F_\theta^s(x)$ to $F_{\theta'}^s(x)$ for $0 \leq s \leq t$. Therefore $d(x, F_\phi^{L\sec(\theta-\theta')}) = d(F_\theta^L(x), F_\phi^{L\sec(\theta-\theta')}(x)) = L\sec(\theta - \theta')\sin(\theta - \phi)$ for any $x$ such that $F_\phi^s(x)$ is contained in the cylinder. The lemma follows by the fact that $|\sin(\theta')| \leq |\theta'|$. $\square$

**Proposition 3.** *If $f$ is a non-increasing positive function $\int_C^{\infty} tf(t)dt = +\infty$ for all $C > 0$ then for almost every $\phi$ and any $L_0$ we have $|\theta - \phi| < f(L)$ where $\phi$ is the direction of a periodic cylinder of volume at least $\sigma$ and length $L > L_0$.*

*Proof.* If $f$ is a non-increasing positive function then $\int_C^{\infty} f(\sqrt{t})dt = \infty$ iff $\int_C^{\infty} tf(t)dt = \infty$. It suffices to prove the discrete version. Observe that $\sum\limits_{i=1}^{\infty} ia_i = \infty$ if and only if $\sum\limits_{i=1}^{\infty} a_{\lfloor\sqrt{i}\rfloor} = \infty$. Indeed, let $M \geq 2$ and notice $\sum\limits_{i=1}^{\infty} M^{2(i-1)}a_{M^i} \leq \sum\limits_{i=1}^{\infty} ia_i \leq \sum\limits_{i=0}^{\infty} M^{2(i+1)}a_{M^i}$, and, on the other hand, $\sum\limits_{i=1}^{\infty} M^{2(i-1)}a_{M^i} \leq \sum\limits_{i=1}^{\infty} a_{\lfloor\sqrt{i}\rfloor} \leq \sum\limits_{i=0}^{\infty} M^{2(i+1)}a_{M^i}$. The proposition now follows by Corollary 4. $\square$

*Proof of Theorem 1.* Because $f$ is non-increasing, if the flow in direction $\phi$ is ergodic it suffices to show that $\liminf\limits_{t\to\infty} f(t)^{-1}d(F_\phi^t(x), x) = 0$ for a positive measure set of $x$. $F_\phi$ is ergodic for Lebesgue almost every $\phi$ by [4]. By Lemma 2 it suffices to show that for any $\epsilon > 0$ there are arbitrarily large $L_0$ such that there is a periodic cylinder of length $L > L_0$ and at least a fixed positive area in direction $\theta$ where $|\phi - \theta| < \frac{\epsilon f(L)}{L}$. This is true by Proposition 3. $\square$

## 1. Acknowledgments

I would like to thank M. Boshernitzan and L. Marchese.

## REFERENCES

[1] Beresnevich, V: Application of the concept of regular systems of points in metric number theory. (Russian) Vestsi Nats. Akad. Navuk Belarusi Ser. Fiz.-Mat. Navuk 140 (2000), no. 1, 35–39.

[2] Boshernitzan, M; Chaika, J: Borel-Cantelli sequences.

[3] Katok, A. B; Zemljakov, A. N: Topological transitivity of billiards in polygons. Mat. Zametki 18 (1975), no. 2, 291–300.

[4] Kerckhoff, S; Masur, H; Smillie, J: Ergodicity of billiard flows and quadratic differentials. Ann. of Math. (2) 124 (1986), no. 2, 293–311.

[5] Marchese, L: Khinchin theorem for interval exchange transformations. J. Mod. Dynam. 5 no. 123-183

[6] Vorobets, Y: Periodic geodesics on translation surfaces. arXiv:math/0307249

*E-mail address*: `jonchaika@math.uchicago.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVENUE, CHICAGO, IL 60637, USA